



# Food insecurity is associated with worse HIV clinical outcomes among women in the United States: Findings from the Women's Interagency HIV Study



Spinelli M<sup>1</sup>, Frongillo EA<sup>2</sup>, Sheira L<sup>3</sup>, Palar K<sup>3</sup>, Tien PC<sup>4</sup>, Wilson T<sup>5</sup>, Merenstein D<sup>6</sup>, Cohen M<sup>7</sup>, Adedimeji A<sup>8</sup>, Wentz E<sup>9</sup>, Adimora A<sup>10</sup>, Ofotokun I<sup>11</sup>, Metsch L<sup>12</sup>, Turan JM<sup>13</sup>, Kushel MB<sup>1</sup>, Weiser SD<sup>3</sup>

<sup>1</sup>Department of Medicine, University of California, San Francisco. <sup>2</sup>Department of Health Promotion, Education, and Behavior, University of South Carolina, Columbia, SC.

<sup>3</sup>Division of HIV, ID and Global Medicine, Department of Medicine, University of California, San Francisco (UCSF), San Francisco, CA.

<sup>4</sup>Department of Medicine, University of California, San Francisco and Medical Service, Department of Veteran Affairs Medical Center, San Francisco, CA.

<sup>5</sup>Department of Community Health Sciences, State University of New York Downstate Medical Center, School of Public Health, Brooklyn, NY.

<sup>6</sup>Department of Family Medicine, Georgetown University Medical Center, Washington, DC. <sup>7</sup>Department of Medicine, Stroger Hospital, Chicago, IL.

<sup>8</sup>Dept. of Epidemiology and Population Health, Albert Einstein College of Medicine, Bronx, NY. <sup>9</sup>Bloomberg School of Public Health, Department of Epidemiology, Johns Hopkins University, Baltimore, MD.

<sup>10</sup>School of Medicine and UNC Gillings School of Global Public Health, University of North Carolina at Chapel Hill, Chapel Hill, NC. <sup>11</sup>School of Medicine, Emory University, Atlanta.

<sup>12</sup>Department of Sociomedical Sciences, Mailman School of Public Health, Columbia, New York, NY. <sup>13</sup>Department of Health Care Organization and Policy, School of Public Health, University of Alabama, Birmingham.

## Introduction

- Food insecurity: limited or uncertain availability of nutritionally adequate, foods or inability to acquire food in socially acceptable ways<sup>1</sup>
- Affects half of HIV-infected patients in the U.S.<sup>2</sup>
- Disproportionately affects women and women-headed households<sup>3</sup>
- Food insecurity associated with higher HIV viral load and lower CD4 counts in populations comprised mostly of men<sup>2,3</sup>
- Adherence proposed as potential mediator between food insecurity and worse HIV clinical outcomes<sup>3</sup>.
- No domestic studies of food insecurity and HIV treatment outcomes specifically among women, none are national<sup>2</sup>

## Results and Outcomes

**Table 1: Characteristics of WIHS participants, n=1,304**

Characteristic	Value
Age, mean (SD)	48.6 (8.7)
Race/ethnicity: African-American, %	68.7
Hispanic, %	16.4
Other, %	3.4
Less than \$500 savings, %	78.2
Having child dependents, %	34.9
Any governmental or non-governmental food aid, %	20.0
Any illicit drug use in last 6 mts, %	23.4
Current alcohol use, %	43.6
Time on ART (years), mean (SD)	9.5 (6.0)
Underweight, BMI<18.5	3.0
Overweight/obese, BMI ≥ 25	72.4
Current non-adherence (<95% of doses), %	17.1
Undetectable viral load, %	58.4
Food insecurity, %	41.9

- **Food insecurity associated with 2.1 times higher HIV-1 viral load (95% CI: 1.1-4.1)**
- Path through adherence accounted for 75.3% of the association between food insecurity and HIV-1 viral load
- **Food insecurity associated with a 43.4 lower mean CD4+ count (-83.1 – -3.6)**
- Adherence accounted for 23.3% of the association

**Table 2: Food insecurity associated with increased HIV-1 viral load and decreased CD4+ count**

	HIV-1 Viral Load Multivariable Adjusted Factor (95% CI)	CD4+ Cell Count Multivariable Adjusted β (95% CI)
Having food insecurity (marginal, low, or very low food security)	2.1 (1.1-4.1)*	-43.4 (-83.1 – -3.6)*
Age per 10 years	0.7 (0.5-1.1)	-
African-American	2.6 (0.8-8.1)	-26.4 (-90.5 – -37.6)
Hispanic white	2.2 (0.6-8.1)	-79.8 (-154.8 – -4.8)*
Other	1.1 (0.6-5.6)	59.4 (-52.7 – 171.5)
At least high school degree	-	-
Less than \$30,000 income	-	-
Less than \$500 savings	-	-48.9 (-97.2 – -0.6)*
Having child dependents	2.7 (1.23-5.8)**	-
Homeless/marginal housing	-	-
Any governmental or non-governmental food aid	-	-94.5 (171.4 – -17.5)*
Any illicit drug use in last 6 mts <sup>2</sup>	3.2 (1.5-7.0)**	-
Current alcohol use	-	9.8 (-36.0 – -55.6)
CD4 nadir	-	-
Time on ART (years)	1.0 (0.9-1.0)	35.1 (26.3 – 44.0)***
Underweight, BMI<18.5	22.9 (3.9-133.8)**	-128.9 (-242.4 – -15.3)*
Overweight/obese, BMI ≥ 25	-	76.6 (32.3 – 120.7)**

\* p<0.05 \*\*p<0.01 \*\*\*p<0.001

## Discussion

- Prior research demonstrates higher HIV-1 viral loads and lower CD4+ counts associated with worse health outcomes
- First to show adherence as mediator for CD4+ count; prior studies show food security to be highly predictive of ART adherence<sup>5</sup>
- Builds on research that suggests women may prioritize their children's health over their own<sup>6</sup>

## Implications

- Further study needed to elucidate the pathways between food insecurity and HIV outcomes
- Assessment of food insecurity could assist programs seeking to bolster ART adherence and improve HIV outcomes
- Comprehensive HIV care that integrates food insecurity interventions may have significant impacts on health of populations living with HIV

## Literature Cited

1. Bickel G, M. Nord, et al. Guide to Measuring Household Food Security, Revised 2000. Alexandria, VA, U.S. Department of Agriculture, Food and Nutrition Service. 2000.
2. Anema A, Weiser SD, Fernandes KA, et al. High prevalence of food insecurity among HIV-infected individuals receiving HAART in a resource-rich setting. *AIDS Care*. 2011;23(2):221-230
3. Weiser SD, Young SL, Cohen CR, et al. Conceptual framework for understanding the bidirectional links between food insecurity and HIV/AIDS. *Am J Clin Nutr*. 2011;94(6):1729S-1739S.
4. Barkan SE, Melnick SL, Preston-Martin S, et al. The Women's Interagency HIV Study. WIHS Collaborative Study Group. *Epidemiology*. 1998;9(2):117-125.
5. Young S, Wheeler AC, McCoy SI, Weiser SD. A review of the role of food insecurity in adherence to care and treatment among adult and pediatric populations living with HIV and AIDS. *AIDS Behav*. 2014;18 Suppl 5:S505-515.
6. McIntyre L, Glanville NT, Raine KD, Dayle JB, Anderson B, Battaglia N. Do low-income lone mothers compromise their nutrition to feed their children? *CMAJ*. 2003;168(6):686-691.

## Acknowledgements

Research funded by NIH R01MH095683 and the U.C.O.P. Global Food Initiative Fellowship. We appreciate the assistance of the Women's Interagency HIV Study (WIHS) for their support.

## Project Aims

**1. Examine association between food insecurity and HIV clinical outcomes in cross-sectional sample from the Women's Interagency HIV Study (WIHS), national sample of HIV-infected women**

**2. Investigate adherence as potential mediator between food insecurity and HIV clinical outcomes**

## Methods

- Sample: Cross-sectional study of 1,304 women throughout U.S as part of Women's Interagency HIV Study (WIHS), national longitudinal study of HIV-infected women<sup>4</sup>
- *Primary Independent Variable:* food insecurity measured using the Household Food Security Survey Module, previously validated in several countries including U.S.
- *Primary outcomes:* continuous HIV-viral load and continuous CD4+ count
- *Covariates:* Demographics, socioeconomic controls, clinical controls, substance use
- *Mediator:* Adherence <95% using a visual analog scale
- *Analysis:* For viral load outcome, used Tobit regression analysis, natural log transformed
- For CD4 outcome, used multivariable linear regression.
- Covariates included with bivariate p<0.15.
- For *mediation analysis*, created alternate model including adherence; measured change in coefficient to determine strength of adherence

